## Maklumat tambahan indikasi

**Tahun 2021** 

## **Products Approved For Additional Indication (DCA 359 – 15 Jun 2021)**

No.	Product [Active Ingredient]	Additional Indication	Marketing Authorization Holder
1.	COMIRNATY Concentrate for Dispersion for Injection [BNT162b2 1 dose (0.3ml) contains 30 µg of modRNA [Single-stranded, 5'-capped mRNA formulated as an RNA-lipid nanoparticle (LNP) of nucleoside- modified mRNA (modRNA)]	INDICATION:  Comirnaty is indicated for active immunisation to prevent COVID-19 caused by SARS-CoV-2 virus, in individuals 12 years of age and older.  The use of this vaccine should be in accordance with official recommendations.  POSOLOGY: Individuals 12 years of age and older  Comirnaty is administered intramuscularly after dilution as a course of 2 doses (0.3 mL each). It is recommended to administer the second dose 3 weeks after the first dose.  There are no data available on the interchangeability of Comirnaty with other COVID-19 vaccines to complete the vaccination course. Individuals who have received 1 dose of Comirnaty should receive a second dose of Comirnaty to complete the vaccination course.	PFIZER (MALAYSIA) SDN. BHD. Level 10 & 11, Wisma Averis, Tower 2, Avenue 5, Bangsar South, No.8, Jalan Kerinchi, 59200 Kuala Lumpur, Wilayah Persekutuan Kuala Lumpur.
		Paediatric population  The safety and efficacy of Comirnaty in children aged less than 12 years of age have not yet been established. Limited data are available.	