

**Year 2007**

**DCA 200 (31/12/2007)**

NO	PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1. 1.1	ACLASTA SOLUTION FOR INFUSION 5mg/100ml  (Zoledronic acid monohydrate)	➤ Treatment of osteoporosis in postmenopausal women to reduce the incidence of hip, vertebral and non-vertebral fractures.	Novartis Corporation (M) Sdn. Bhd. Level 15, CREST, 3 Two Square No. 2, Jalan 19/1 46300 Petaling Jaya Selangor
2. 2.1	NEXAVAR FILM-COATED TABLETS 200mg  (Sorafenib)	➤ Treatment of patients with hepatocellular carcinoma	Bayer Schering Pharma Bayer Co. (M) Sdn Bhd TI-14 Jaya 33 No. 3 Jalan Semangat Seksyen 13 46200 Petaling Jaya Selangor
3. 3.1 3.2	Aricept 5mg Tablet  Aricept 10mg Tablet  (Donepezil hydrochloride)	➤ Treatment of mild to moderate dementia in Alzheimer's disease, as well as in patients with severe Alzheimer's disease.	Eisai (Malaysia) Sdn. Bhd. Lot 6.1, 6 <sup>th</sup> Floor Menara Lien Hoe No. 8, Persiaran Tropicana 47410 Petaling Jaya Selangor
4. 4.1 4.2 4.3	AVELOX TABLET  AVELOX INFUSION SOLUTION - 250ml VIAL  AVELOX INFUSION SOLUTION - 250ml FLEXIBAG  (Moxifloxacin)	Avelox is indicated for the treatment of adults (≥18 years of age) with the following bacterial infections caused by susceptible strains:  ➤ Complicated intra-abdominal infections including polymicrobial infections such as abscesses.	Zuellig Pharma Sdn Bhd Level 3A, Block B No. 10, Jalan Bersatu 13/4 46200 Petaling Jaya Selangor

**DCA 199 (04/12/2007)**

NO	PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1. 1.1	ARIXTRA 12.5mg/ml SOLUTION FOR INJECTION  (Fondaparinux Sodium)	➤ Treatment of acute Deep Vein Thrombosis(DVT) ➤ Treatment of acute Pulmonary Embolism(PE)	Glaxosmithkline Pharmaceutical Sdn. Bhd. 8th Floor, Menara Lien Hoe No.8, Persiaran Tropicana 47410 Petaling Jaya  Selangor
2. 2.1	ACTONEL 35mg TABLET  (Risedronate Sodium)	➤ Treatment of osteoporosis in men at high risk of fracture.	Sanofi-Aventis(M) Sdn. Bhd. 8th Floor, PNB Damansara No. 19, Lorong Dungun Damansara Heights 50490 Kuala Lumpur
3. 3.1	REMICADE POWDER FOR CONCENTRATE FOR SOLUTION	➤ Treatment of severe, active Crohn's disease in paediatric patients aged 6 to	Schering-Plough Sdn. Bhd. Level 3-1 CP Tower

CONCENTRATE FOR SOLUTION  
FOR INFUSION

(Infliximab)

disease, in paediatric patients aged 6 to 17 years, who have not responded to conventional therapy including a corticosteroid, an immunomodulator and primary nutrition therapy; or who are intolerant to or have contraindications for such therapies. Remicade has been studied only in combination with conventional immunosuppressive therapy.

Level 17, C1 Tower,  
No. 11, Jalan 16/11,  
Pusat Dagang Seksyen 16,  
46350 Petaling Jaya  
Selangor

**DCA 197 (27/9/2007)**

NO		PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1	ARIXTRA 2.5MG/0.5ML SOLUTION FOR INJECTION  (Fondaparinux Sodium)	<p>1. Treatment of unstable angina or non-ST segment elevation myocardial infarction (UA/NSTEMI) in patients for whom urgent (120mins) invasive management (PCI) is not indicated.</p> <p>1. Treatment of ST segment elevation myocardial infarction (STEMI) in patients who are managed with thrombolytics or who initially are to receive no other form of reperfusion therapy.</p>	Glaxosmithkline Pharmaceutical Sdn. Bhd. 8th Floor, Menara Lien Hoe No.8, Persiaran Tropicana 47410 Petaling Jaya Selangor
2.	2.1 2.2	CYMBALTA CAPSULE 30MG  CYMBALTA CAPSULE 30MG  (Duloxetine Hydrochloride)	Treatment of generalized anxiety disorder.	Eli Lilly (Malaysia) Sdn. Bhd. Unit 18.1, Level 18, Cp Tower No. 11, Jalan 16/11 Pusat Dagang Seksyen 16 46350 Petaling Jaya Selangor
3.	3.1 3.2 3.3	TARCEVA FILM-COATED TABLET 25 MG  TARCEVA FILM-COATED TABLET 100 MG  TARCEVA FILM-COATED TABLET 150 MG  (Erlotinib Hydrochloride)	<p><u>Pancreatic Cancer:</u></p> <p>Tarceva in combination with gemcitabine is indicated for the first-line treatment of patients with locally advanced, unresectable or metastatic pancreatic cancer.</p>	Roche (M) Sdn. Bhd. 14th. Floor, West Block Wisma Selangor Dredging, 142, Jalan Ampang 50450 Kuala Lumpur
4.	4.1 4.2 4.3	ZOCOR TABLETS 10MG  ZOCOR TABLETS 20MG  ZOCOR TABLETS 40MG  (Simvastatin)	<p><u>Paediatric Patients with Heterozygous Familial Hypercholesterolemia</u></p> <p>Zocor is indicated as an adjunct to diet to reduce total-C, LDL-C, TG and Apo-B levels in adolescent boys and girls who are at least</p>	Merck Sharp & Dohme(I.A.) Corp. Letter Box 1502 15th Floor, Menara Merais No. 1, Jalan 19/3 46300 Petaling Jaya

			one year post-menarche, 10-17 years of age, with heterozygous familial hypercholesterolemia (HeFH).	Selangor
5.	5.1	FORTEO 250 UG SOLUTION FOR INJECTION  (Teriparatide)	To increase bone mass in men with primary or hypogonadal osteoporosis who are at high risk for fracture.	Eli Lilly (Malaysia) Sdn. Bhd. Unit 18.1, Level 18, Cp Tower No. 11, Jalan 16/11 Pusat Dagang Seksyen 16 46350 Petaling Jaya Selangor

**DCA 196 (30/8/2007)**

NO		PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1	PEG-INTRON POWDER FOR INJECTION 50MCG	PEG-INTRON is indicated for the treatment of chronic Hepatitis-C and Chronic Hepatitis-B. Patients must be 18 years of age or older and have compensated liver disease.	Schering-Plough Level 3-1 CP Tower No 11, Jalan 16/1 Pusat Dagang Seksyen 16 46350 Petaling Jaya Selangor Darul Ehsan
	1.2	PEG-INTRON POWDER FOR INJECTION 80MCG		
	1.3	PEG-INTRON POWDER FOR INJECTION 100MCG		
	1.4	PEG-INTRON POWDER FOR INJECTION 120MCG		
	1.5	PEG-INTRON POWDER FOR INJECTION 150MCG		
2.	2.1	PEG-INTRON REDIPEN 50MCG		
	2.2	PEG-INTRON REDIPEN 80MCG		
	2.3	PEG-INTRON REDIPEN 100MCG		
	2.4	PEG-INTRON REDIPEN 120MCG		
	2.5	PEG-INTRON REDIPEN 150MCG		

**DCA 195 (07/8/2007)**

NO		PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1	NICORETTE 2MG CHEWING GUM	As an aid to reduce smoking, prior to stopping completely (Nicotine Assisted Reduction Stop [NARS]).	Pfizer (Malaysia) Sdn. Bhd. Level 3 & 4, Bangunan Palm Grove No. 14, Jalan Glenmarie, Section U1 40150 Shah Alam Jaya Selangor
	1.2	NICORETTE 4MG CHEWING GUM  (Nicotine)		
2.	2.1	AVANDAMET TABLETS 1MG/ 500MG	Avandamet is indicated as an adjunct to diet and exercise for the treatment of type 2	Glaxosmithkline Pharmaceutical Sdn. Bhd.

	2.2	AVANDAMET TABLETS 2MG/ 500MG	diabetes mellitus (non-insulin dependent diabetes mellitus). Avandamet is indicated as initial therapy, to improve glycemic control in patients with type 2 diabetes mellitus who are not adequately controlled on diet and exercise alone.	8th Floor, Menara Lien Hoe No.8, Persiaran Tropicana 47410 Petaling Jaya Selangor
	2.3	AVANDAMET TABLETS		
	2.4	4MG/ 500MG		
	2.5	AVANDAMET TABLETS 2MG/ 1000MG		
		AVANDAMET TABLETS 4MG/ 1000MG  (Rosiglitazone Maleate / Metformin Hydrochloride)		
3.	3.1	TRAVATAN 0.004% OPHTHALMIC SOLUTION  (Travoprost)	TRAVATAN® OPTHALMIC SOLUTION is indicated for the reduction of elevated intraocular pressure in patients with open-angle glaucoma or ocular hypertension.	Alcon Laboratories (Malaysia) Sdn. Bhd. No. 1001A & 1001D, 9th Floor Wisma Glomac 3 Kompleks Kelana Centre Point No. 3, Jalan SS7/19, Kelana Jaya 47301 Petaling Jaya Selangor
4.	4.1	TAXOTERE 20MG/0.5ML VIAL	<u>Head and Neck Cancer</u> Taxotere in combination with cisplatin and 5-fluoracil is indicated for the induction treatment of patients with inoperable locally advanced squamous cell carcinoma of the head and neck.	Sanofi-Aventis (Malaysia) Sdn. Bhd. 8th Floor, PNB Damansara No. 19, Lorong Dungun Damansara Heights 50490 Kuala Lumpur
	4.2	TAXOTERE 80MG/2.05ML VIAL  (Docetaxel)		
5.	5.1	BETAFERON INJECTION  (Recombinant Interferon Beta-1B)	Betaferon is indicated for:  1. the treatment of patients with a single clinical event suggestive of multiple sclerosis and at least two clinically silent magnetic resonance imaging (MRI) lesion characteristic of multiple sclerosis, if alternative diagnoses been excluded.	Zuellig Pharma Sdn Bhd Level 3a, Blok B 10, Jalan Bersatu 13/4 46200 Petaling Jaya Selangor

**DCA 194 (21/6/2007)**

NO	PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
----	--------------------------------	-----------------------	--------------------------------------

1.	1.1	EZETROL 10MG (Ezetimibe)	<u>Primary Hypercholesterolemia</u> Ezetrol, administered in combination with fenofibrate, is indicated as adjunctive therapy to diet for the reduction of elevated total-C, LDL-C, Apo B and non-HDL-C in patients with mixed hyperlipidaemia.	Merck Sharp & Dohme(I.A.) Corp. Letter Box 1502 15th Floor, Menara Merais No. 1, Jalan 19/3 46300 Petaling Jaya Selangor
2.	2.1	PLAVIX TABLET 75MG (Clopidogrel Hydrogen Sulphate)	Clopidogrel is indicated for the prevention of atherothrombotic events in:  1. Patients suffering from acute coronary syndrome:  - ST segment elevation acute myocardial infarction, in combination with acetyl salicylic acid (ASA) in medically treated patients eligible for thrombolytic therapy.	Sanofi-Aventis (Malaysia) Sdn. Bhd. 8th Floor, PNB Damansara No. 19, Lorong Dungun Damansara Heights 50490 Kuala Lumpur.

**DCA 193 (24/5/2007)**

NO		PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1 1.2 1.3	MAXIPIME FOR INJECTION 0.5G/VIAL MAXIPIME FOR INJECTION 1.0G MAXIPIME INJECTION 2G/VIAL (Cefepime)	Maxipime is also indicated for surgical prophylaxis in patients undergoing intra-abdominal surgery. In this indication, it is essential that metronidazole also be administered.	Bristol-Meyers Squibb (M) Sdn Bhd 17th. Floor, Menara Lien Hoe 8, Persiaran Tropicana 47410 Petaling Jaya Selangor.
2.	2.1	NEXIUM IV 40 MG (Esomeprazole)	Nexium for injection and infusion is indicated for gastric antisecretory treatment when the oral route is not possible, such as:  1. Healing of gastric ulcers associated with NSAID therapy 2. Prevention of gastric and duodenal ulcers associated with NSAID therapy, in patients at risk.	Astrazeneca Sdn. Bhd. Ground Floor, Wisma Prima 17, Jalan Semantan Satu 50490 Kuala Lumpur.
3.	3.1	THYMOGLOBULINE 25MG (Rabbit anti human thymocyte immunoglobulin 5mg/1ml)	Prophylaxis of acute and chronic graft versus host disease (GvHD), after haematopoietic stem cell transplantation.	Diethelm Malaysia Sdn Bhd 74, Jalan Universiti 46200 Petaling Jaya Selangor

**DCA 192 (3/5/2007)**

NO		PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1	FEMARA 2.5MG FILM-COATED TABLET  (Letrozole)	Pre-operative therapy in postmenopausal women with localized hormone receptor positive breast cancer, to allow subsequent breast-conserving surgery in women not originally considered candidates for this type of surgery. Subsequent treatment after surgery should be in accordance with standard of care.	NOVARTIS CORPORATION (M) SDN BHD LOT 9, JALAN 26/1, SEKSYEN 26, KAWASAN PERINDUSTRIAN HICOM 40400 SHAH ALAM, SELANGOR
2.	2.1 2.2	LUCRIN DEPOT 3.75MG INJECTION LUCRIN DEPOT 11.25MG INJECTION  (Leuprolide Acetate)	Leuprolide Acetate is indicated for the treatment of breast cancer in pre-menopausal women in whom hormone therapy is specified.	ABBOTT LABORATORIES (M) SDN BHD NO 22, JALAN PEMAJU U1/15, SEKSYEN U1, HICOM-GLENMARIE INDUSTRIAL PARK, 40150 SHAH ALAM, SELANGOR.

**DCA 191 (29/3/2007)**

NO		PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1 1.2	SEROXAT CR 12.5 MG SEROXAT CR 25 MG  (Paroxetine hydrochloride)	<u>Premenstrual Dysphoric Disorder (PMDD)</u> Seroxat CR tablets are indicated for the treatment of premenstrual dysphoric disorder (PMDD).  Paroxetine has not been evaluated beyond 3 menstrual cycles in controlled clinical trials for the treatment of PMDD.	GLAXOSMITHKLINE PHARMACEUTICAL SDN BHD 8TH FLOOR, MENARA LIEN HOE, NO. 8 PERSIARAN TROPICANA, 47410 PETALING JAYA, SELANGOR.
2.	2.1	HYCAMTIN INJECTION 4 MG  (Topotecan hydrochloride 4mg)	Hycamtin in combination with cisplatin is indicated for the treatment of patients with histologically confirmed Stage IV-B, recurrent, or persistent carcinoma of the cervix, which is not amenable to curative treatment with surgery and/or radiation therapy.	GLAXOSMITHKLINE PHARMACEUTICAL SDN BHD 8TH FLOOR, MENARA LIEN HOE, NO. 8 PERSIARAN TROPICANA, 47410 PETALING JAYA, SELANGOR

3.	3.1 3.2 3.3 3.4	SEROQUEL TABLET 25MG SEROQUEL TABLET 100MG SEROQUEL TABLET 200MG SEROQUEL TABLET 300MG  (Quetiapine Fumarate)	Treatment of depressive episodes associated with bipolar disorder  The efficacy of Seroquel was established in two identical 8-week randomized, placebo-controlled double-blind clinical studies that included either bipolar I or II patients with and without rapid cycling courses. Effectiveness has not been systematically evaluated in clinical trials for more than 8 weeks.	ASTRAZENECA SDN. BHD. GROUND FLOOR WISMA PRIMA 17 JALAN SRI SEMANTAN SATU DAMANSARA HEIGHTS 50490 KUALA LUMPUR.
4.	4.1	REMICADE POWDER FOR CONCENTRATE FOR SOLUTION FOR INFUSION  (Infliximab 100mg/vial)	<u>Ulcerative colitis</u>  REMICADE is indicated for reducing signs and symptoms, inducing and maintaining clinical remission and mucosal healing, and eliminating corticosteroid use in patients with moderately to severely active ulcerative colitis who have had an inadequate response to conventional therapy.	SCHERING-PLOUGH SDN BHD LEVEL 3-1, CP TOWER NO. 11, JALAN 16/11 PUSAT DAGANG SEKSYEN 16 46350 PETALING JAYA, SELANGOR.

**DCA 190 (22/2/2007)**

NO		PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1 1.2 1.3	NOVOSEVEN 60 KIU / VIAL (1.2MG / VIAL) POWDER FOR INJECTION  NOVOSEVEN 120 KIU/ VIAL (2.4MG/ VIAL) POWDER FOR INJECTION  NOVOSEVEN 240 KIU/ VIAL (4.8MG/ VIAL) POWDER FOR INJECTION  (Activated recombinant human coagulation Factor VIIa (rFVIIa))	Treatment of bleeding episodes and for prevention of bleeding in those undergoing surgery or invasive procedures :  <ul style="list-style-type: none"> <li>▪ in patients with congenital Factor VII (FVII) deficiency.</li> </ul>	NOVO NORDISK PHARMA (M) SDN. BHD. 11.02, 11TH FLOOR, MENARA YAYASAN SELANGOR NO. 18A, JALAN PERSIARAN BARAT 46000 PETALING JAYA SELANGOR
2.	2.1 2.2 2.3 2.4	EXELON 1.5MG CAPSULES EXELON 3.0 MG CAPSULES EXELON 4.5 MG CAPSULES EXELON 6.0 MG	<ul style="list-style-type: none"> <li>▪ Treatment of patients with mild to moderately severe dementia associated with Parkinson's disease.</li> </ul>	NOVARTIS CORPORATION (M) SDN. BHD. LOT 9, JALAN 26/1 SEKSYEN 26 KAWASAN PERINDUSTRIAN

	2.5	CAPSULES EXELON ORAL SOLUTION 2MG/ML (120ML)		HICOM 40000 SHAH ALAM SELANGOR.
	2.6	EXELON ORAL SOLUTION 2MG/ ML (50ML)		
		(Rivastigmine hydrogen tartrate)		
3.	3.1	NEXIUM 20 MG TABLET	<ul style="list-style-type: none"> <li>■ Treatment of Zollinger Ellison Syndrome.</li> </ul>	ASTRAZENECA SDN. BHD. GROUND FLOOR WISMA PRIMA 17 JALAN SRI SEMANTAN SATU DAMANSARA HEIGHTS 50490 KUALA LUMPUR.
	3.2	NEXIUM 40 MG TABLET  (Esomeprazole magnesium trihydrate)		
4.	4.1	PARIET 10 MG	<ul style="list-style-type: none"> <li>■ Symptomatic treatment of moderate to very severe gastro-esophageal reflux disease (Symptomatic GERD).</li> </ul>	EISAI (M) SDN. BHD. LOT 6.1, 6TH FLOOR MENARA LIEN HOE NO. 8, PERSIARAN TROPICANA 47410 PETALING JAYA SELANGOR.
	4.2	PARIET 20 MG  (Rabeprazole sodium)		

**DCA 189 (25/1/2007)**

NO		PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1	ZYVOX TABLET 400MG	Zyvox formulations are indicated in the treatment of complicated skin and skin structure infections, including diabetic foot infections, without concomitant osteomyelitis, caused by Staphylococcus aureus, (methicilin-susceptible and resistant strains), Streptococcus pyogenes, or Streptococcus agalactiae. Zyvox has not been studied in the treatment of decubitus ulcers. Combination therapy may be clinically indicated if the documented or presumptive pathogens include Gram-negative organisms.	Pfizer (Malaysia) Sdn. Bhd. Level 3 & 4, Bangunan Palm Grove, No. 14, Jalan Glenmarie (Persiaran Kerjaya), Section U1, 40150 Shah Alam, Selangor.
	1.2	ZYVOX TABLET 600MG		
	1.3	ZYVOX GRANULES FOR ORAL SUSPENSION 20MG/ML		
	1.4	ZYVOX INFUSION BAGS 200MG/100ML		
	1.5	ZYVOX INFUSION BAGS 600MG/300ML  (Linezolid)		
2.	2.1	INVANZ INJECTION (Ertapenem sodium)	Invanz is indicated in adults for the prophylaxis of surgical site infection following elective colorectal surgery.	Merk Sharp & Dohme (I.A.) Corp. Malaysia Branch,



				Letter Box 1502, 15th Floor Menara Merais, 1, Jalan 19/3, 46300 Petaling Jaya, Selangor.
3.	3.1 3.2 3.3	COVERSYL 2MG TABLET COVERSYL 4MG TABLET COVERSYL 8MG TABLET  (Perindopril tertbutylamine)	<u>Stable Coronary Artery Disease</u>  Reduction of risk of cardiac events in patients with a history of myocardial infarction and/or revascularization.	Servier (M) Sdn. Bhd. 1301, Level 13, Uptown 2, Damansara Utama, 47400 Petaling Jaya, Selangor.
4.	4.1 4.2 4.3	AVELOX TABLET AVELOX SOLUTION VIAL- INFUSION 250ML AVELOX SOLUTION FLEXIBAG- INFUSION 250ML  (Moxifloxacin)	Avelox is indicated for the treatment of adults ( $\geq 18$ years of age) with complicated skin and skin structure infections caused by susceptible strains.	Zuellig Pharma Sdn. Bhd. Level 3A, Block B, No. 10, Jalan Bersatu 13/4, 46200 Petaling Jaya, Selangor.
5.	5.1 5.2	TAMIFLU CAPSULES 75MG TAMIFLU ORAL SUSPENSION 12MG/ML  (Oseltamivir phosphate)	Tamiflu is indicated for post exposure prophylaxis of influenza in adults and children aged 1 year and above.	Roche (M) Sdn. Bhd. 14th Floor, West Block, Wisma Selangor Dredging, 142, Jalan Ampang, 50450 Kuala Lumpur.
6.	6.1	HUMIRA SOLUTION FOR INJECTION  (Recombinant human somatropin)	<u>Rrheumatoid Arthritis</u> Humira is indicated for reducing signs and symptoms, inducing major clinical response and clinical remission, inhibiting the progression of structural damage and improving physical function in adult patients with moderately to severely active rheumatoid arthritis. Humira can be used alone or in combination with methotrexate or other disease modifying anti-rheumatic drugs (DMARDs).  <u>Psoriatic Arthritis</u> Humira is indicated for reducing the signs and symptoms of active arthritis in patients with psoriatic arthritis. Humira can be used alone or in combination with disease modifying anti-rheumatic drugs (DMARDs).  <u>Ankylosing Spondylitis</u> Humira is indicated for reducing signs and symptoms in patients with active ankylosing spondylitis.	Abbott Laboratories (M) Sdn. Bhd. No. 22, Jalan Pemaju U1/15 Section U1 HICOM-Glenmarie Industrial Park 40150 Shah Alam Selangor.
7.	7.1	NORDITROPIN SIMPLEXX 5 MG / 1.5ML INJECTION	Growth retardation in prepubertal children due to chronic renal disease.	Novo Nordisk Pharma (M) Sdn. Bhd.

7.2	<p>NORDITROPIN NORDILET 5 MG / 1.5 ML INJECTION</p>	<p>Growth disturbance (current height standard deviation score (SDS) &lt; -2.5 and parental adjusted height SDS &lt; -1) in short children born small for gestational age (SGA), with a birth weight and/or length below -2 standard deviation (SD), who failed to show catch-up growth (height velocity standard deviation score (HV SDS) &lt; 0 during the last year) by 4 years of age or later.</p>	<p>11.02, 11th Floor Menara Yayasan Selangor No. 18A, Jalan Persiaran Barat 46000 Petaling Jaya Selangor.</p>
7.3	<p>NORDITROPIN NORDILET 10 MG / 1.5 ML INJECTION</p> <p>(Recombinant Human Somatropin)</p>		